



GASTROENTEROLOGY SPECIALISTS, INC.

10210 East 91st Street South
Tulsa, OK 74133
(918) 940-8500

Medication Instructions

- * **High Blood Pressure Patients:** Take your morning dose of blood pressure medication.
- * **Diabetic Patients on Insulin:** Take 1/2 dose of your PM insulin the day before your procedure.
- * **Diabetic Patients:** Do not take any insulin or diabetic medication by mouth the day of your procedure **UNLESS** your blood sugar is higher than 300g/dL. If this occurs, take insulin according to your sliding scale.
- * After consulting with your physician, **STOP TAKING:**

Blood Thinners		
⇒ If you take 81mg Aspirin for heart protection, DO NOT STOP IT.		
7 Days before procedure	5 Days before procedure	2 Days before procedure
<ul style="list-style-type: none"> • Plavix (Clopidogrel) • Aggrenox • Effient (prasugrel) 	<ul style="list-style-type: none"> • Brilinta (Ticagrelor) • Coumadin (warfarin) • Fish oil 	<ul style="list-style-type: none"> • Eliquis • Pradaxa • Xarelto

If taking below medications for Diabetic management and/or Weight Loss STOP:	
7 Days before procedure	Day of procedure
<ul style="list-style-type: none"> • Ozempic (Semaglutide) • Wegovy (Semaglutide) • Mounjaro (Tirzepatide) • Trulicity (Dulaglutide) • Rybelsus (Semaglutide) • Bydureon (Exenatide) 	<ul style="list-style-type: none"> • Byetta (Exanatide) • Victoza (Liraglutide) • Saxenda (Liraglutide)

****Due to potential anesthesia complications, your procedure may be canceled if you do not stop medications per recommendation****

3 Days before procedure
<ul style="list-style-type: none"> • Iron and Bulk Laxatives • Metamucil • Effersyllium • Avoid high-fiber foods

EGD Preparation Instructions

- ⇒ Please do not smoke the day of your procedure.
- ⇒ **High Blood Pressure Patients:** Take your morning dose of blood pressure medication.

To avoid possible procedure cancellation:

1. Do not eat or drink anything before your procedure as explained below
2. Arrive for your procedure **45 minutes** before your scheduled procedure time
3. Bring a responsible adult who can remain at GSI throughout your entire procedure
4. Any female who has had a menstrual period within the last year will be required to take a urine pregnancy test upon arrival. Please do not urinate in the lobby.

Please follow if taking medications listed in the box below for Diabetes and/or Weight Loss

7 Days before procedure STOP:	Day of procedure STOP:
<ul style="list-style-type: none">• Ozempic (Semaglutide)• Wegovy (Semaglutide)• Mounjaro (Tirzepatide)• Trulicity (Dulaglutide)• Rybelsus (Semaglutide)• Bydureon (Exenatide)	<ul style="list-style-type: none">• Byetta (Exanatide)• Victoza (Liraglutide)• Saxenda (Liraglutide)

Day before your procedure

- * **When you wake up, begin a clear liquid diet. STOP any liquids 8 hours before your procedure arrival time.**
 - Do not eat any solid foods or dairy products or drink alcohol.**
 - Clear liquids include: water, soda, sports drinks, clear broth or bouillon, black coffee, black tea, popsicles, plain Jell-O, and clear fruit juices without pulp (white grape, apple juice)
- !! IMPORTANT !! NOTHING in your mouth 8 Hours before your procedure arrival time. DO NOT:**
 - Eat food, mints, gum, or candy
 - Drink anything, including water
 - Chew tobacco or nicotine

Please follow if not taking medications listed in the box above

- * **Diabetic Patients taking insulin:** Take 1/2 dose of your PM insulin the day before your procedure.

On the day of your procedure

- * **After midnight, DO NOT:**
 - Eat food, mints, or candy
 - Drink anything, including water
 - Chew tobacco or nicotine
- * You may brush your teeth.
- * **Diabetic Patients taking insulin:** Do not take any insulin or diabetic medication by mouth the day of your procedure **UNLESS** your blood sugar is higher than 300g/dL. If this occurs, take insulin according to your sliding scale.



GASTROENTEROLOGY SPECIALISTS, INC.

10210 East 91st Street South
Tulsa, OK 74133
(918) 940-8500
(918) 940-8399 -Fax

Home Care Instructions After Your Procedure

During your procedure you will be given medication that may affect your memory of the day's events. Other normal side effects of the medicine may include confusion, irritability, drowsiness or a drunken-like state. Resting at home is the best way to recover.

Your speech, posture, hand control, and ability to work may be affected for 24 hours. A responsible person needs to drive you home after the procedure, and someone should stay with you for your safety until the effects of the medication have subsided.

- * Do not make any important decisions or sign important documents for 24 hours.
- * Do not drive or operate any machinery for 24 hours.

After your procedure, resume your usual diet unless instructed otherwise by your physician.

Do not drink alcoholic beverages for 24 hours.

If you take blood thinning medication you should ask your physician when it is safe to resume.

If you take prescription medicines on a regular basis, ask your physician which ones you should take after your procedure.

Do not take any muscle relaxants, sedatives, hypnotics or mood altering medicines for 24 hours unless ordered by your physician. If you have pain, you may take over the counter medicines such as Tylenol.

Call your physician if you have any of the following symptoms after your procedure:	
<ul style="list-style-type: none">• Blue Skin• Unable to Arouse• Bleeding	<ul style="list-style-type: none">• Difficulty Breathing• Pain Not Relieved by Medicine• Fever or Chills

GSI Phone Number: (918) 940-8500

www.gsitulsa.com